

Claims:

1. A method of representing a three dimensional tooth object for use in determine an orthodontic setup of the teeth, comprising:

displaying data of a three-dimensional object for lower molars;
5 selecting groove and cusp points from the displayed object; and
defining from the selected groove and cusp points line segments representative of the orientation of the tooth object.

2. The method of claim 1 wherein:

the selecting includes selecting a distal-lingual groove point (DLG.Pt), a distal-lingual cusp point
10 (DLC.Pt), a mesial-lingual cusp point (MLC.Pt), a facial cusp point (FC.Pt), and a mesial-facial
cusp point (MFC.Pt) from the displayed object; and
the defining includes defining from points DLC.Pt and MLC.Pt a labial cusp line segment (LC.Li),
from points FC.Pt and MFC.Pt a facial cusp line segment (FC.Li), from points DLG.Pt and
MFC.Pt a line segment Stolar.Li.

15 3. The method of claim 1 further comprising calculating the occlusion of upper molars based
on the defined line segments.

4. The method of claim 1 further comprising:

defining an occlusal table line segment (OT.Li) perpendicular to LC.Li, coincident with the buccal
groove, and connecting FC.Li to LC.Li, representing a mid-developmental lobe plane (MDL.Pl).

20 5. The method of claim 4 further comprising:

defining the initial mesial-distal positioning of an appliance on the tooth in relation to the mid-
developmental lobe plane (MDL.Pl).

6. The method of claim 1 wherein:

the selecting includes:

25 selecting a mesial-lingual cusp point (MLC.Pt), a distal-lingual cusp point (DLC.Pt), a mesial
facial cusp point (MFC.Pt), and a distal-facial cusp point (DFC.Pt), and
selecting mesial and distal marginal ridges as points MMR.Pt and DMR.Pt, respectively; and

the defining includes:

defining from points MLC.Pt and DLC.Pt a labial cusp line segment (LC.Li), from points MFC.Pt
30 and DFC.Pt a facial cusp line segment (FC.Li),
defining a marginal ridge line segment (MR.Li) as that through points MMR.Pt and DMR.Pt, and
defining from points MMR.Pt and DLC.Pt a line segment Stolar.Li.

- 22 -

REPLACED BY
ART 34 AMDT

7. The method of claim 7 further comprising:
calculating the occlusion with lower molars using the line Stolar Li.

8. The method of claim 7 further comprising:
selecting points MFC.Pt and MLC.Pt to define mesial cusp line segment (MC.Li);
5 defining occlusal table line segment (OT.Li) to be parallel to line MC.Li, coincident with the buccal
groove, and connecting FC.Li to LC.Li; and
defining with line OT.Li the mid-developmental lobe plane (MDL.Pl).

9. The method of claim 8 further comprising defining the initial mesial-distal positioning of
an appliance on the tooth in relation to the mid-developmental lobe plane (MDL.Pl).

10. A method of forming a digital image of a tooth comprising:
forming a 3-D tooth object of the crown of a tooth by scanning the tooth or a model thereof; and
adding tooth root data to the tooth object from a separate file or object that contains three dimensional
data of one or more individual tooth roots.

11. The method of claim 10 wherein the tooth root data is produced by scans of the patient's
15 individual anatomy.

12. The method of claim 10 wherein the teeth root data is retrieved from library files or records
that contain three-dimensional shape data of anatomically representative tooth roots.

13. The method of any of claims 10-12 further comprising:
modifying the tooth root data from information from the tooth crown scans.

20 14. The method of any of claims 10-13 further comprising:
joining the crown and root data to form complete tooth objects by aligning the crown long axis and
the root long axis from the respective data.

15. The method of claim 14 further comprising:
setting the alignment to a selected or calculated relative torque angle.

25 16. The method of any of claims 10-15 further comprising:
using root image data that represent the tooth root at less than full scale relative to that of the crown;
incrementally increasing in size the tooth root data until any part of the root touches a corresponding
point in the crown;

- 23 -

when this contact occurs, fixing the expansion of the root at the contact point and proportionately scaling the rate of expansion from the contact point until the root has been expanded and the rims of the crown and root fully coincide.

5 17. The method of calculating an orthodontic setup of a patient's teeth including calculating the rotation of individual teeth to produce an occlusion.

18. The method of calculating an orthodontic setup of a patient's teeth including calculating a symmetrical arch by superimposing images of contralateral teeth in MDL.Pl cross-sectional views with their PAW.Pts coincident and rotating images of the teeth about the PAW.Pt until predetermined occlusal angles of the teeth are at an average value.

10 19. The method of claim 18 further comprising extruding or intruding the teeth in a direction normal to the occlusal plane until the buccal cusp points of the teeth are coincident with the occlusal plane.

20. The method of claim 19 further comprising defining the horizontal distance from the buccal cusp point (BC.Pt) to PAW.Pt as the prominence line (PR.Li), and establishing said line to an average value by adjusting the BC.Pts.

15 21. The method of calculating an orthodontic setup of a patient's teeth including calculating a symmetrical arch comprising fixing FIE.Pts of the lower incisors to align the lower incisal edges to a best fit buccal cusp curve to enable placement of upper incisors, setting the BC.Pt of teeth lower incisors to an average value from FIE.Pt along MDL.Pl, finding the intersection of MDL.Pl and midline from the mandibular trough for lower central incisors and calculating the line segment distance from BC.Pt along
20 MDL.Pl to intersection for both teeth.

22. The method of claim 21 further comprising calculating from the intersections, a respective distance equal to said line segments facially along midline, averaging points at the measured distances on midline to define thereby the origin of an anterior bezier handle, and defining the mesial facial cusps of lower molars as posterior bezier handles.

25 23. A method of determining a dental arch for arrangement of a patient's teeth comprising: displaying dental anatomy of a patient on a display for adjustment by an operator; and representing a dental arch on the display with a bezier curve having anterior bezier and posterior bezier handles by which the operator can alter the shape of the arch on the display.

30 24. The method of claim 22 or claim 23 further comprising moving the handles by an operator to adjust the curve and setting the curve as a default best fit buccal cusp curve.

- 24 -

REPLACED BY
ART 34 AMDT

25. A method of calculating an orthodontic setup of teeth comprising:
providing an operator with an interface on which to select teeth for extraction;
calculating the setup eliminating therefrom the teeth selected by an operator on the interface for
extraction.

5 26. The method of claim 25 further comprising:
providing an operator with the option of selecting teeth on different arches or different sides of
an arch differently;
calculating the setup separately for each arch or side of an arch to provide for asymmetric
extraction.

10 27. A method of providing a custom orthodontic appliance for treatment of a patient, the method
comprising:

maintaining a database, accessible by a computer, and containing data related to each of a plurality of
orthodontic practitioners;

storing in the database, information identifying the practitioners and information relating to treatment
plan options preferred by the respective practitioners;

15 receiving information from an orthodontic practitioner of the plurality for providing a custom
orthodontic appliance for a patient;

in response to the information from the practitioner, determining parameters for the configuration of
a custom orthodontic appliance for the patient based at least in part on the stored options preferred
by the orthodontic practitioner; and

20 providing an orthodontic appliance design for the patient having a configuration that includes the
determined parameters.

28. The method of claim 27 further comprising:

receiving information from the orthodontic practitioner of a treatment plan option for treatment of the
patient; and

25 determining at least one of the parameters based at least in part on the information of the treatment
plan option received from the orthodontic practitioner.

29. The method of claim 27 further comprising:

receiving information from the orthodontic practitioner of a treatment plan option for treatment of the
patient; and

30 determining parameters addressed by the information of the treatment plan option received from the
orthodontic practitioner in accordance with that information and determining parameters not so
addressed in accordance with information stored in the database.

- 25 -

30. A method of determining an arrangement of the teeth of a patient to achieve by orthodontic treatment, comprising:

scanning shapes of the patient's teeth to generate one or more data files thereof;
deriving an arrangement of the patient's teeth from the data with the aid of a computer;
5 taking steps to reduce the amount of the data prior to deriving the arrangement.

31. The method of claim 30 wherein:

said steps include pruning the files to minimize unnecessary information the deriving of the arrangement.

32. The method of claim 30 wherein:

10 the scanning includes scanning a model or impression of the teeth;
said steps include manicuring or trimming the model or impression before scanning.

33. A method of further comprising:

scanning shapes of a plurality of teeth of a patient and generating data of the plurality; then
15 processing the generated data automatically using feature recognition software to isolate tooth objects;
then
displaying the automatically processed generated data to an operator to resolve ambiguities or errors
or to verify the isolated objects.

34. The method of claim 33 further comprising:

manipulating the displayed data to repair the isolated objects to remove artifacts:

20 35. The method of claim 34 wherein:

the manipulating includes modifying tooth crown shapes in the data to eliminate interferences.

36. The method of claim 34 wherein:

the manipulating is performed on a computer display by an operator with the aid of software tools.

37. The method of claim 34 wherein:

25 the manipulating includes selecting, then elevating, lowering or smoothing areas that could cause poor
jig adaptation.

38. The method of claim 33 further comprising:

thinning data of unnecessary resolution.

39. A method of determining an orthodontic setup of the teeth, comprising:

30 defining a mandibular computer model of an arrangement of a patient's mandibular teeth on a
computer; and

ART 34 AMDT

- 26 -

defining a plane of occlusion from the mandibular computer model by identifying three points on different ones of the teeth on the model.

40. The method of claim 39 wherein:

the three points include the mesial buccal cusps of the first mandibular molars and the incisal tip of one of the mandibular centrals.

41. A method of determining an arch shaped arrangement for an orthodontic setup of the teeth, comprising:

generating on a computer a mandibular model of the patient's teeth that includes gingival or jaw shape data;

generating a section through the model through the gingiva or jaw; and

selecting points defining facial and lingual boundaries and defining thereby a mandibular trough of the patient.

42. The method of claim 41 further comprising:

displaying the mandibular trough on a computer display;

calculating a smooth symmetrical bezier curve through the trough between corresponding labial and lingual points;

the curve having handles at the midline and at the distal ends thereof to allow an operator to modify the shape of the bezier curve.

43. A method of determining an orthodontic setup of the teeth, comprising:

producing data representing each of a plurality of a patient's teeth;

defining for each tooth a mid-developmental lobe plane.

44. The method of claim 43 wherein the defining includes:

defining for the tooth, a mid-developmental lobe axis (MDL.Ax) by selection of its extents incisally (IMDL.Pt) and gingivally (GMDL.Pt);

defining through MDL.Ax and perpendicular to a mesial-distal axis of the tooth, a mid developmental lobe plane.

45. The method of claim 41 further comprising:

define a crown long axis for the tooth in the mid developmental lobe plane.

46. A method of representing a three dimensional tooth object for use in determine an orthodontic

setup of the teeth, comprising:

arranging tooth objects at angles of the teeth for each tooth in their own XYZ coordinate systems, in which Z is perpendicular to an occlusal plane and X is a mesial-distal axis for the tooth;

- 27 -

REF
ART 34 AMDT

wherein tooth torque angle is set in a plane perpendicular to the X axis, tip angle is set as the angle of the plane relative to the X axis.

47. The method of claim 46 further comprising:
defining a set-up by arranging the coordinate systems of the teeth.

5 48. The method of claim 46 further comprising:
defining a set-up by arranging the coordinate systems of the teeth with the coordinate systems rotated relative to each other to a value determinative of tooth rotation.